










The Distance to the Vela Supernova Remnant

Alexandra N. Cha¹, Kenneth R. Sembach¹, Anthony C. Danks²

Received _____; accepted _____

arXiv:astro-ph/9902230v2 17 Feb 1999

¹Department of Physics & Astronomy, The Johns Hopkins University, Baltimore, MD 21218

²Raytheon STX, Goddard Space Flight Center, Code 683, Greenbelt, Maryland 20771

ABSTRACT

We have obtained high resolution Ca II and Na I absorption line spectra toward 68 OB stars in the direction of the Vela Supernova Remnant. The stars lie at distances of 190 – 2800 pc as determined by Hipparcos and spectroscopic parallax estimations. The presence of high velocity absorption attributable to the remnant along some of the sight lines constrains the remnant distance to 250 ± 30 pc. This distance is consistent with several recent investigations that suggest that the canonical remnant distance of 500 pc is too large.

Subject headings: line: profiles – ISM: clouds – ISM: individual (Vela Supernova Remnant) – ISM: supernova remnants

1. Introduction

Supernova remnant (SNR) studies provide valuable information about the amount of energy released in supernova explosions, the effects of blast waves on the structure and physical conditions in the interstellar medium (ISM), shock physics, and dust grain destruction. Distance is a fundamental quantity that often factors into such studies. Knowledge of SNR distances is crucial for determining and justifying calculations of remnant ages (Wallerstein & Silk 1971), predicting the explosion energies of the progenitor supernovae, and quantifying temperatures behind shock fronts (Jenkins & Wallerstein 1995).

The Vela SNR is one of the best studied SNRs in the sky. An uncertain distance of 500 pc (Milne 1968) is often quoted. Recent studies of the conditions within the remnant suggest that a smaller value is more appropriate. We have undertaken an observing program to determine the distance to the Vela SNR directly by spectroscopic means. We present these results in this *Letter* and provide additional results pertaining to time variability in the high velocity absorption in a companion paper (Cha & Sembach 1999, hereafter Paper II).

2. Data

We obtained high-resolution Ca II K (3933.663 Å) and Na I D₂,D₁ (5889.951, 5895.924 Å) spectra of 68 OB stars in the direction of the Vela SNR using the Coudé Echelle Spectrograph on the Coudé Auxiliary Telescope at the European Southern Observatory in 1993, 1994, and 1996. The instrumental setups are described in Paper II along with the data reduction procedures (see also Sembach et al. 1993). The fully reduced data have typical signal-to-noise ratios in excess of 100 and 2-pixel resolutions of $\lambda/\Delta\lambda \approx 75,000$,

or $\approx 4 \text{ km s}^{-1}$, as determined from the widths of Th-Ar lamp spectra. The spectra are presented in Paper II.

The stars observed rotate rapidly and serve as suitable background sources for studies of narrow interstellar (or remnant) absorption lines. The locations of the stars are shown in Figure 1 along with a ROSAT 0.75 keV X-ray contour (Snowden et al. 1995) arising from the extreme edge of the adiabatically expanding shock encircling the remnant.

3. Previous Distance Estimates

The distance to the Vela SNR has been questioned ever since Milne (1968) proposed a distance of $500 \pm 100 \text{ pc}$. Milne arrived at this value by comparing the observed angular diameters of the Vela SNR, Cygnus Loop, and IC 443, and assuming that all three had similar linear diameters. Comparing the Cygnus Loop ($\phi = 170'$, $d = 770 \text{ pc}$; Minkowski 1958; Harris 1962) to Vela, ($\phi = 4^\circ$), yielded $d_{\text{Vela}} \approx 540 \text{ pc}$. A similar comparison of Vela to IC 443 ($\phi = 45'$, $d = 2 \text{ kpc}$ – Milne’s “probable distance”) resulted in $d_{\text{Vela}} \approx 375 \text{ pc}$. The Vela SNR distance of $500 \pm 100 \text{ pc}$ thus arose from a rough average of these distances. Kristian (1970) summarizes five early distance measurements, similarly concluding that $d_{\text{Vela}} = 500 \text{ pc}$, but with a factor of two uncertainty.

A compilation of recent distance estimates to the Vela SNR can be found in Table 1. The methods included a repeat of Milne’s calculations using an updated angular diameter of the Vela SNR (Oberlack et al. 1994). That work, which also nicely recounts previous distance determinations, and suggests $d_{\text{Vela}} = 230 \pm_{115}^{230} \text{ pc}$. A second distance estimate involved an analysis of the proper motion of the Vela pulsar by Ögelman et al. (1989), who found that if the distance to the pulsar were $d_{\text{Vela}} = 290 \pm 80 \text{ pc}$, then the projected velocity of the pulsar matched the pulsar velocity derived from interstellar scintillation data (Cordes

1986).

As early as 1971, Wallerstein & Silk (1971) recommended a distance near 250 pc to unify age estimates of the remnant derived from the observed rate of pulsar deceleration with estimates based on the shock velocity and angular size of the remnant. More recently, Jenkins & Wallerstein (1995) quantified the total energy of the Vela SNR and found that a distance of 500 pc resulted in an energy that was an order of magnitude larger than the expected $\approx 10^{51}$ ergs for Type II supernovae (Woosley & Weaver 1986; Arnett et al. 1989). A factor of 2 smaller distance lowered the energy by a factor of 8 to a more representative value. Jenkins & Wallerstein (1995) also noted the high correlation between the theoretically calculated temperature behind the shock wave and the observational temperature estimates of the X-ray emitting regions of Vela (Kahn et al. 1985) if the adopted distance to Vela were ≈ 250 pc.

4. A Precise Distance to the Vela SNR

We have analyzed high-resolution Ca II absorption spectra of 68 stars in the direction of the Vela SNR ($l \approx 264^\circ$, $b \approx -3^\circ$). Many of the spectra reveal Ca II and Na I absorption at velocities well beyond those expected for gas in the local interstellar medium (LISM), which has $V_{\text{LSR}}(\text{Ca II}) \approx 0.9 \pm 9.4 \text{ km s}^{-1}$ (Vallerga et al. 1993). In the direction of Vela, we find no ISM absorption toward δ Vel ($d \approx 20$ pc). Crawford (1991) found interstellar Ca II at velocities of -6.6 and -1.6 km s^{-1} toward κ Vel ($d \approx 100$ pc). These velocities are typical of many southern hemisphere (Sco-Cen-Vel) sight lines extending 100–200 pc from the Sun; the southern hemisphere LISM components span the -20 to $+6 \text{ km s}^{-1}$ velocity range (Crawford 1991; Welsh et al. 1994; Génova et al. 1997). In contrast to these relatively low LISM mean velocities, Dubner et al. (1998) reported the presence of H I 21cm emission from gas about the center of the SNR accelerated to velocities of $|V_{\text{LSR}}| \approx 30 \text{ km s}^{-1}$ by the

expansion of the Vela supernova shock.

To determine the distance to the Vela SNR, we decomposed each Ca II profile into Gaussian components and identified those components that appear at velocities indicative of the remnant. In Figure 2 we plot the central velocities of each component as a function of background star distance. To illustrate the characteristically low velocity foreground gas at $100 \text{ pc} < d < 200 \text{ pc}$, we also plot data for stars in the Sco-Cen region (Crawford 1991). We estimated stellar distances and errors using two methods: trigonometric parallaxes based upon Hipparcos parallax measurements and spectroscopic parallaxes based upon photometric colors and spectral types reported in the literature (see Paper II). There was good agreement between the two methods, but whenever possible, we adopted the more accurate Hipparcos trigonometric parallax values.

Given the systematically low velocities of LISM material and the higher velocity H I emission seen by Dubner et al. (1998) toward Vela, we adopted a lower limit of $\pm 25 \text{ km s}^{-1}$ for absorption that we considered to be associated with the remnant. A much more conservative limit of $\pm 50 \text{ km s}^{-1}$ could also be adopted, and for the sake of erring on the side of caution we will discuss our distance estimate with this higher velocity cutoff when appropriate. These velocities are highlighted by dashed and solid vertical lines, respectively, in Figure 2. Beyond 250 pc, all of our Ca II spectra but one (HD 74662) show absorption components at $|V_{\text{LSR}}| > 25 \text{ km s}^{-1}$, and beyond 390 pc absorption lines with $|V_{\text{LSR}}| > 50 \text{ km s}^{-1}$ appear in many of the spectra.

We show the Ca II K spectrum of two stars at 250 pc, HD 70309 ($d_{\text{Hipp}} = 250 \pm 35 \text{ pc}$) and HD 71459 ($d_{\text{Hipp}} = 250 \pm 30 \text{ pc}$), in Figure 3. Three absorption components at $\langle V_{\text{LSR}} \rangle = -28, -13, \text{ and } +1 \text{ km s}^{-1}$ are readily discernible in the spectrum of HD 70309. The -13 and $+1 \text{ km s}^{-1}$ components are also present in the spectrum of nearer stars and probably arise in the LISM at a distance of 40–200 pc. We believe the -28 km s^{-1} component reveals

gas associated with the front side of the remnant. The spectrum of HD 71459 also reveals higher velocity Ca II absorption. Figure 3 shows that there is an absorption line with $\langle V_{\text{LSR}} \rangle = +30 \text{ km s}^{-1}$, indicating the presence of remnant gas moving away from us. The two sight lines are separated by 6.2° , or about 27 pc at a distance of 250 pc. This is roughly the diameter of the remnant at this distance. Both sight lines pass near the edge of the X-ray emission line boundary of the remnant, so the total space velocities of the remnant gas along the sight lines could be considerably higher than the measured velocities.

We now consider a simple model to determine what the deprojected space velocities for these two sight lines might be. We assume that the remnant is a spherically expanding shell centered in the X-ray contour, with approximate coordinates $\alpha_{2000} = 08^h 40^m$ and $\delta_{2000} = -45^\circ 00'$.³ Applying a spherical coordinate system allows the total space velocity of the gas to be resolved into vector components, including the observed velocity, which is the projection of the total velocity onto the line of sight. We assign a right handed coordinate system with the origin at the center of the remnant and the following axes: the x-axis extends from the observer to the remnant, the y-axis is right ascension, and the z-axis is declination. Converting from spherical to Cartesian coordinates, we find the space velocity, v , knowing the observed velocity (x-projection), v_{obs} , to be $v_{\text{obs}} = v \sin \theta \cos \phi$. From Figure 1, we estimate $\theta \approx 45^\circ$ for HD 71459 and $\theta \approx 130^\circ$ for HD 70309. Determining ϕ would require knowing the distance to clouds on the spherical remnant. A robust lower limit on the total velocity is obtained by setting $\phi = 0^\circ$ (i.e., $v > v_{\text{obs}} \csc \theta$).

Applying the projection effects to the absorption components of the two stars at 250 pc, we find that the observed velocity of the component at -28 km s^{-1} toward HD 70309

³We note that the Ca II data is not described well by a simplistic spherical model (see Wallerstein et al. 1980), so we use it only as a first order approximation to illustrate the general effects of projection.

converts to a minimum space velocity of -40 km s^{-1} , and the observed velocity of the component at $+30 \text{ km s}^{-1}$ toward HD 71459 corresponds to a minimum space velocity of $+39 \text{ km s}^{-1}$. These space velocities are indeed much higher than those expected for gas associated with the LISM, lending support to our claim that observed LSR velocities beyond $\pm 25 \text{ km s}^{-1}$ toward these stars arise within the remnant.

Higher velocity absorption is also seen along some sight lines toward Vela. Figure 3 contains the Ca II spectrum of HD 72350, a B4 IV star at $d = 390 \pm 100 \text{ pc}$, as determined using the standard reddening relation, $A_v = 3.1 E(B-V)$, with the UBV photometry of Schild et al. (1983). Here, absorption components are observed at intermediate velocities, $V_{\text{LSR}} = -20$ and $+38 \text{ km s}^{-1}$, as well as at high velocities, $V_{\text{LSR}} = +114$ and $+122 \text{ km s}^{-1}$. The spectrum of another star at a similar distance, HD 74319 ($d_{\text{Hipp}} = 400 \pm 90 \text{ pc}$), also reveals high velocity absorption components, though at negative velocities, $V_{\text{LSR}} = -61$ and -74 km s^{-1} . Both stars are located near the center of the X-ray shell (Figure 1). The observed absorption velocities along these sight lines are probably close to the actual space velocities of the shocked regions they probe.

The distribution of sight lines with widely varying absorption signatures is testimony to the patchiness of the remnant region. The observed distant and nearby stars are randomly projected onto the remnant, revealing no obvious bias in their spatial distribution. We calculated the two-sided Kolmogorov-Smirnov (K-S) statistic on the spectral components velocities for the stars at $245 \text{ pc} < d < 385 \text{ pc}$ and $385 \text{ pc} < d < 555 \text{ pc}$ and found it to be, $D=0.14$, with a significance level of 0.40, implying that the two groups of stars are not from significantly different populations.

Assigning gas with observed velocities of $|V_{\text{LSR}}| > 25 \text{ km s}^{-1}$ to the Vela SNR results in an SNR distance of $250 \pm 30 \text{ pc}$. An upper limit of $\sim 390 \text{ pc}$ is allowed if only $|V_{\text{LSR}}| > 50 \text{ km s}^{-1}$ components are retained in the analysis. In light of the LISM data, we believe that

the upper limit is overly pessimistic, and therefore we recommend a distance of 250 ± 30 pc be used for future studies of the remnant.

We report and recalculate some general characteristics of the Vela SNR in Table 2 using a distance of 250 ± 30 pc. The recalculations of the various attributes of the remnant follow the prescriptions outlined in the previously published work described in §3.

5. Supporting Absorption Line Data

We have reviewed published ultraviolet absorption line data to test our observationally determined distance to the Vela SNR. Choosing components with $|V_{\text{LSR}}| > 25 \text{ km s}^{-1}$ in the International Ultraviolet Explorer (IUE) data reported by Jenkins et al. (1984) yields an upper limit of ~ 400 pc to the distance, based upon an uncertain detection of C IV at -33 km s^{-1} toward HD 72555 ($d = 390\pm90$ pc), and the detection of a -125 km s^{-1} absorption component in low ionization lines (Fe II, Si II) toward HD 74436 ($d = 410\pm230$ pc). Copernicus data for stars in the direction of the Vela SNR (Wallerstein et al. 1980) yield an upper limit of 480 ± 120 pc as constrained by the sight line to HD 72127A. The distance obtained using the Copernicus data set is almost a factor of two higher than our measurement, but this may be attributed to the limited sample of 14 stars, which results in a much coarser sampling of distances. The IUE sample was more extensive (45 sight lines) and yields an upper bound consistent with our distance estimate. The major advantage of our spectroscopic investigation over previous absorption line studies of the region is the combination of high signal to noise and high spectral resolution, resulting in a high sensitivity to weak lines that may not have been observable in the Copernicus and IUE data.

6. Conclusions

The analysis of a significant sample of O and B stars in the direction of the Vela SNR has enabled a direct distance measurement of 250 ± 30 pc, with a conservative upper limit of 390 ± 100 pc. This measurement is consistent with recent arguments that favor a smaller distance to Vela.

We appreciate comments on this work from our colleagues at the Johns Hopkins University. KRS and ANC acknowledge support from NASA Long Term Space Astrophysics Grant NAG5-3485 and grant GO-06413-95A from the Space Telescope Science Institute, which is operated by AURA under NASA contract NAS5-26555.

REFERENCES

- Arnett, W.D., Bahcall, J.N., Kirshner, R.P., & Woosley, S.E. 1989, ARA&A, 27, 629
- Aschenbach, B. 1993, AdSpR, 131, 45
- Aschenbach, B. 1998, Nature, 396, 141
- Cha, A.N., Sembach, K.R., & Danks, A.C. 1999, ApJS, submitted (Paper II)
- Cordes, J.M. 1986, ApJ, 311, 183
- Crawford, I.A. 1991, A&A, 247, 183
- Dubner, G.M., Green, A.J., Goss, W.M., Bock, D.C.-J., & Giacani, E. 1998, AJ, 116, 813
- Génova, R., Beckman, J., Bowyer, S., & Spicer, T. 1997, ApJ, 484, 761
- Harris, D.E. 1962, ApJ, 135, 661
- Jenkins, E. & Wallerstein, G. 1995, ApJ, 440, 227
- Jenkins, E., Wallerstein, G., & Silk, J. 1984, ApJ, 278, 649
- Kahn, S. M., Gorenstein, P., Harnden, F. R., & Seward, F. D. 1985, ApJ, 299, 821
- Kristian, J. 1970, ApJ, 162, L103
- Milne, D.K. 1968, Australian J. Phys., 21, 201
- Minkowski, R. 1958, Rev. Mod. Phys., 30, 1048
- Oberlack, U., Diehl, R., Montmerle, T., Prantzos, N., & von Ballmoos, P. 1994, ApJS, 92, 433
- Ögelman, H., Koch-Miramond, L., & Aurière, M. 1989, ApJ, 342, L83
- Prentice, A. & ter Haar, D. 1969, Nature, 222, 964
- Schild, R.E., Garrison, R.F., & Hiltner, W.A. 1983, ApJS, 51,321

- Snowden, S.L., Freyberg, M.J., Plucinsky, P.P., Schmitt, J.H.M.M., Truemper, J., et al. 1995, ApJ, 454, 643
- Vallerga, J. V., Vedder, P. W., Craig, N., & Welsh, B. Y. 1993, ApJ, 411, 729
- Wallerstein, G. & Silk, J. 1971, ApJ, 170, 289
- Wallerstein, G., Silk, J., & Jenkins, E.B. 1980, ApJ, 240, 834
- Welsh, B.Y., Craig, N., Vedder, P.W., & Vallerga, J.V. 1994, ApJ, 437, 638
- Woosley, S.E., & Weaver, T.A. 1986, ARA&A, 24, 205

Figure Captions

Fig. 1.— Positions of the sample stars in the sky. The stars discussed in §4 are identified. An “X” marks the location of the Vela pulsar. The ROSAT X-ray contour represents 4×10^{-4} counts s^{-1} arcmin^{-2} .

Fig. 2.— Central velocities (diamonds) of the observed Ca II absorption components embedded with horizontal ticks revealing the Doppler spread parameters versus the distance of the corresponding star. Crosses indicate Ca II velocity data by Crawford (1991).

Fig. 3.— Ca II K absorption spectra towards HD 70309, HD 71459, HD 72350, and HD 74319. Tick marks identify absorption by gas within the SNR.

TABLE 1
VELA SNR DISTANCE MEASUREMENTS

Reference	Distance	Comments
Milne (1968)	500 ± 100 pc	Angular diameter comparison with Cygnus Loop and IC 443
Prentice & ter Haar (1969)	410 pc	Pulsar dispersion measure determination (see Davies 1969).
Davies (1969)	600 pc	Pulsar dispersion measure determination (see Prentice & ter Haar 1969).
Wallerstein & Silk (1971)	~ 250 pc	Required for consistency in age estimates.
Ögelman et al. (1989)	290 ± 80 pc	Required for consistency with transverse motion of Vela pulsar.
Oberlack et al. (1994)	$230 \pm_{15}^{230}$ pc	Repeat of Milne's (1968) approach with new angular diameters.
Jenkins & Wallerstein (1995)	~ 250 pc	Required for an acceptable value of the total energy of the SNR.
This paper	250 ± 30 pc	Spectroscopic identification of high velocity SNR gas.

TABLE 2
ATTRIBUTES OF THE VELA SNR AT 250 ± 30 PC

Attribute	Value	Comment/Reference
Diameter	32 ± 4 pc	Using angular size of 7.3° (Aschenbach 1993).
Age of Remnant	11,400 yr	Pulsar spin-down age (Reichley et al. 1970).
	$13,000 \text{ yr} < t < 16,000 \text{ yr}$	Shock velocity considerations (Wallerstein & Silk 1971).
Energy of Remnant	$1\text{--}2 \times 10^{51}$ ergs	See Jenkins & Wallerstein (1995).
Temperature Behind Blast	4.5×10^6 K	See Jenkins & Wallerstein (1995).